



## Dr. Satyapriya Sahoo

**Qualification:** MBBS, MD(Paediatrics), DrNB(Neonatology)

**Department:** Pediatrics

**Designation:** Consultant Neonatologist and Pediatrician

**Languages:** Telugu, Hindi, English, Odia

**Experience:** 7 yrs

### About the Doctor

---

Dr. Satyapriya Sahoo is a Consultant Neonatologist and Paediatrician working at Yashoda Hospitals, Malakpet.

### Educational Qualifications

---

- April 2022 to April 2025: DrNB Neonatology, Fernandez Hospital, Hyderabad
- 2018 to 2021: Senior Residency, AIIMS, Bhubaneswar
- 2018: M.D. (Pediatrics), SVPPGIP and SCB Medical College, Cuttack, Odisha, Utkal University
- 2009: B.S., S.C.B. Medical College, Cuttack, Utkal University

### Experience

---

- July 2018 to July 2021: Senior Resident, Department of Pediatrics & Neonatology AIIMS Bhubaneswar.
- July 2021 to April 2022: Junior Consultant, Department of Pediatrics and Neonatology at Sum Ultimate Hospital, Bhubaneswar

## Services Offered

---

- PICC, UAC/UVC catheters
- Thoracocentesis and intercostal drain insertion
- Arterial cannulation
- Endotracheal intubation
- Peritoneal dialysis catheterization
- Lumbar puncture
- Surfactant- LISA
- Initiation and maintenance of mechanical ventilation (conventional, CPAP)
- HFOV with INO, NIV, HFNC
- USG Guided Procedures in Emergency and ICU based Management
- Therapeutic hypothermia.
- USG Guided Procedures in Emergency and ICU based Management
- Therapeutic hypothermia.
- Neonatal functional ECHO.
- Neonatal EEG interpretation

## Awards and Achievements

---

- Advanced NRP certification
- Certification for Neurodevelopmental assessment (Bayley)

## Professional Membership

---

- IMA (Indian Medical Association)
- IAP (Indian Academy of Paediatrics)
- NNF (National Neonatology Forum)

## Research & Publications

---

- Clinical profile of down syndrome in children less than 14 years in a tertiary care hospital- A prospective cohort study
- The profile of serum and urinary sodium levels and its correlation with growth outcomes among preterm newborns less than 32 weeks - A prospective cohort study